

Photoproduction and ultra-peripheral collisions with Pythia 8

POETIC 8

Ilkka Helenius

March 20th, 2018

Tübingen University
Institute for Theoretical Physics

In collaboration with
Torbjörn Sjöstrand

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



LUND
UNIVERSITY



Why study photoproduction?

- Monte-Carlo event generators essential to study the potential of future experiments
- Photo-nuclear processes can be used to probe the structure of nucleons (nuclear PDFs)
- ⇒ Connection between EIC and ultra-peripheral heavy ion collisions at the LHC

Outline

1. Photoproduction in PYTHIA 8
2. Comparisons to HERA data
3. Ultraperipheral heavy-ion collisions
4. Summary & Outlook

Photoproduction in PYTHIA 8

- A general-purpose Monte-Carlo event generator
- Current version 8.230, next release within a few weeks
- Main focus has been in pp, now extensions to ee, ep, pA, AA

Team:

- Torbjörn Sjöstrand Lund University
- Christian Bierlich Lund University
- Nishita Desai CNRS-Universite de Montpellier
- Nadine Fischer Monash University
- Ilkka Helenius Tübingen University
- Philip Ilten University of Birmingham
- Leif Lönnblad Lund University
- Stephen Mrenna Fermi National Accelerator Laboratory
- Stefan Prestel Fermi National Accelerator Laboratory
- Christine O. Rasmussen Lund University
- Peter Skands Monash University

Event generation in PYTHIA 8

1. Hard process generation

- Generate according to LO partonic cross section and PDFs (or feed in processes from external matrix element generator)

2. Parton showers

- Generate Initial and Final State Radiation (ISR & FSR) according to DGLAP evolution equations

3. Multiparton interactions (MPIs)

- Use regularized QCD $2 \rightarrow 2$ cross sections finite also at $p_T \rightarrow 0$

4. Add beam remnants

- Minimal number of partons to conserve colour and flavour
- Fix momenta so that total momentum is conserved

5. Hadronization

- Using Lund string model with color reconnection
- Decays into stable hadrons

Photoproduction in ep

Photoproduction: Small photon virtuality $Q_\gamma^2 \lesssim 1 \text{ GeV}^2$ (cf. DIS)

- Factorize the flux of photons from the hard scattering (Weizsäcker-Williams)


$$f_\gamma^l(x_\gamma) = \frac{\alpha_{em}}{2\pi} \frac{(1 + (1 - x_\gamma)^2)}{x_\gamma} \log \left[\frac{Q_{\max}^2}{Q_{\min}^2(x_\gamma)} \right]$$

- Direct processes**

- Photon initiator of the hard process
- No MPIs but FSR and ISR for hadron



- Resolved processes**

- Photon fluctuates into a hadronic state
- Partonic structure described with PDFs
- FSR and ISR for both sides, also MPIs



PDFs for resolved photons

Obtained through global DGLAP analysis (LEP data mainly)




- Some differences between analyses, especially for gluon
⇒ Theoretical uncertainty for resolved processes
- CJKL used as a default in PYTHIA 8, others via LHAPDF5 but only for hard-process generation

MPIs in PYTHIA 8

- Probability for MPIs from $2 \rightarrow 2$ QCD processes
- Partonic cross section diverges at $p_T \rightarrow 0$
⇒ Regulate the divergence with screening parameter p_{T0}


$$\frac{d\sigma^{2 \rightarrow 2}}{dp_T^2} \propto \frac{\alpha_s(p_T^2)}{p_T^4} \rightarrow \frac{\alpha_s(p_{T0}^2 + p_T^2)}{(p_{T0}^2 + p_T^2)^2}$$

- pp: Power-law in \sqrt{s}
 $p_{T0}(\sqrt{s}) = p_{T0}^{\text{ref}}(\sqrt{s}/7 \text{ TeV})^\alpha$
 $p_{T0}^{\text{ref}} = 2.28 \text{ GeV}/c, \alpha = 0.215$
(Monash tune)
- $\gamma\gamma$: Logarithmic in \sqrt{s}
 $p_{T0}(\sqrt{s}) = p_{T0}^{\text{ref}} + \alpha \log(\sqrt{s}/100 \text{ GeV})$
 $p_{T0}^{\text{ref}} = 1.52 \text{ GeV}/c, \alpha = 0.413$
(I.H., T. Sjöstrand, *in prep.*)
- Parametrization for γp ?



Comparisons to HERA data

Charged particle p_T spectra in ep collisions at HERA



[H1: Eur.Phys.J. C10 (1999) 363-372]


H1 measurement

- $E_p = 820 \text{ GeV}$, $E_e = 27.5 \text{ GeV}$
- $\langle W_{\gamma p} \rangle \approx 200 \text{ GeV}$
- $Q_\gamma^2 < 0.01 \text{ GeV}^2$

Comparison to PYTHIA 8

- Resolved contribution dominates
 - Good agreement with the data using $p_{T0}^{\text{ref}} = 3.00 \text{ GeV}/c$
- \Rightarrow MPI probability between pp and $\gamma\gamma$

Charged particle p_T spectra in ep collisions at HERA



[H1: Eur.Phys.J. C10 (1999) 363-372]


H1 measurement

- $E_p = 820 \text{ GeV}, E_e = 27.5 \text{ GeV}$
- $\langle W_{\gamma p} \rangle \approx 200 \text{ GeV}$
- $Q_\gamma^2 < 0.01 \text{ GeV}^2$

Comparison to PYTHIA 8

- Resolved contribution dominates
 - Good agreement with the data using $p_{T0}^{\text{ref}} = 3.00 \text{ GeV}/c$
- ⇒ MPI probability between pp and $\gamma\gamma$

Charged particle p_T spectra in ep collisions at HERA



H1 measurement

- $E_p = 820$ GeV, $E_e = 27.5$ GeV
- $\langle W_{\gamma p} \rangle \approx 200$ GeV
- $Q_\gamma^2 < 0.01$ GeV²

Comparison to PYTHIA 8

- Resolved contribution dominates
 - Good agreement with the data using $p_{T0}^{\text{ref}} = 3.00$ GeV/c
- \Rightarrow MPI probability between pp and $\gamma\gamma$

Dijet photoproduction in ep collisions at HERA

ZEUS dijet measurement

- $Q_\gamma^2 < 1.0 \text{ GeV}^2$
- $134 < W_{\gamma p} < 277 \text{ GeV}$
- $E_T^{\text{jet1}} > 14 \text{ GeV}$,
 $E_T^{\text{jet2}} > 11 \text{ GeV}$
- $-1 < \eta^{\text{jet1,2}} < 2.4$


Different contributions

- Define

$$x_\gamma^{\text{obs}} = \frac{E_T^{\text{jet1}} e^{\eta^{\text{jet1}}} + E_T^{\text{jet2}} e^{\eta^{\text{jet2}}}}{2yE_e}$$

to discriminate direct and resolved processes
($=x$ in γ at LO parton level)


- At high- x_γ^{obs} direct processes dominate



[ZEUS: Eur.Phys.J. C23 (2002) 615-631]

Dijet in ep collisions at HERA


Pseudorapidity dependence of dijets [Eur.Phys.J. C23 (2002) 615-631]



- Simulations tend to overshoot the dijet data by $\sim 10\%$
- $\sim 10\%$ uncertainty from photon PDFs for $x_\gamma^{\text{obs}} < 0.75$

Ultraperipheral heavy-ion collisions


Motivation: Nuclear parton distribution functions (nPDFs)




- ⇒ Large uncertainties especially for gluon nPDFs
- ⇒ Uncertainty in the pQCD baseline for heavy-ion physics at the LHC

Data available for nPDF fits

- Fixed-target (ν)DIS and DY
 - Pions in dAu at RHIC
 - Dijets in pPb at the LHC
 - EW bosons at the LHC
- ⇒ Limited kinematic reach



Ultra-peripheral heavy-ion collisions



- Large impact parameter $b \Rightarrow$ No strong interaction
- EM-field of nuclei described with quasi-real photons (EPA)
 \Rightarrow Flux of photons with low virtuality (= Photoproduction)
 - **Photon-photon** (dileptons, light-by-light)
 \Rightarrow Useful to calibrate the photon flux
 - **Photon-nucleus** (dijets, incl. hadrons, heavy flavours, ...)
 \Rightarrow Can be used to probe nuclear PDFs

Proposed by M. Strikman, R. Vogt and S. White [PRL 96 (2006) 082001]

Photon-photon interactions

Photon flux from nuclei in impact-parameter b space

- Obtained by a Fourier transformation of the time-dependent EM-field

$$x_\gamma f_\gamma^A(x_\gamma, b) = \frac{\alpha_{\text{EM}} Z^2}{\pi^2} \left[\frac{x_\gamma m}{\hbar c} K_1 \left(\frac{x_\gamma b m}{\hbar c} \right) \right]^2$$



where Z is nuclear charge, m (per-nucleon) mass and K_1 modified Bessel function [Jackson, Classical Electrodyn., 2nd ed.]

Effective photon-photon luminosity

- Need to reject events with hadronic interactions
 - Reject events based on hard-sphere approximation
 - ⇒ Possible to set up in PYTHIA 8
 - Use hadronic interaction probabilities based on nuclear overlap, e.g. STARLIGHT [Comput.Phys.Commun. 212 (2017) 258-268]

High-mass dimuons in ultraperipheral Pb+Pb at the LHC

$$\text{Pb+Pb} \rightarrow \mu^+ + \mu^- + \text{Pb}^* + \text{Pb}^*$$



- Data well described by STARLIGHT MC
- ⇒ Confirms EPA for Pb+Pb at the LHC

- PYTHIA hard-sphere flux agrees with STARLIGHT
- Small difference at high- W from nuclear density (\sim high- x_γ)

Photon-nucleus interactions

Flux for photon-nucleus interactions

- Integrate over $b > 2R_A$ to reject hadronic interactions

$$x_\gamma f_\gamma^A(x_\gamma) = \frac{2\alpha_{\text{EM}}Z^2}{\pi} \left[\xi K_1(\xi)K_0(\xi) - \frac{\xi^2}{2} (K_1^2(\xi) - K_0^2(\xi)) \right],$$

where $\xi = 2R_A x_\gamma m / \hbar c$



- Maximum $W_{\gamma\text{Pb}} \approx 2\sqrt{s}$ in HERA

Photo-nuclear dijet production

- ATLAS analysis [ATLAS-CONF-2017-011]
anti- k_T , $R = 0.4$, $p_T^{\text{lead}} > 20$ GeV, $p_T^{\text{jets}} > 15$ GeV, $|\eta| < 4.4$
- Event-level variables:

$$m_{\text{jets}} = \sqrt{(\sum_i E_i)^2 - |\sum_i \vec{p}_i|^2}, \quad H_T = \sum_i p_{Ti}$$
$$y_{\text{jets}} = \frac{1}{2} \log \left(\frac{\sum_i E_i + \sum_i p_{zi}}{\sum_i E_i - \sum_i p_{zi}} \right), \quad x_A = \frac{m_{\text{jets}}}{\sqrt{s}} e^{-y_{\text{jets}}}$$


Differential photo-nuclear dijet distributions (Preliminary)



EPPS16 will be implemented to the next PYTHIA release

- The expected nPDF features visible in x_A : shadowing, etc..
- Nuclear modifications only in hard-process cross sections

Expected potential of the dijet data



Dominant contribution


- Large x_A : resolved
- Small x_A : direct

Expected statistical error

- Assume $L = 1 \text{ nb}^{-1}$ for the measurement
- Clearly smaller than nPDF uncertainty

⇒ Potential to provide further constraints for nPDFs down to $x \approx 10^{-3}$

Dijet η distribution



Dijet kinematics

- Due to soft γ spektrum jets asymmetrically distributed in η
- No need to push for large η to gain sensitivity to small x

Quantifying the impact of the data to nPDFs requires

- Finalized data
- NLO calculation for photoproduction of dijets
- Accurate description of photon flux from nuclei

Summary & Outlook

Photoproduction implemented into PYTHIA 8

- Automatic mixing of direct and resolved processes
- Full parton-level evolution (parton showers, MPIs)
- Agreement with HERA data, support for MPIs
- Can simulate UPCs by using heavy-ion specific photon flux (though not yet with nuclear target but with nPDFs)

Ultra-peripheral heavy-ion collisions

- Use dilepton production to calibrate the photon flux
- Can study photo-nuclear processes with LHC before EIC
- Dijets provide nPDF constraints down to $x \sim 10^{-3}$
- Number of potential observables, increased low- x reach at lower p_T

Things to do

- Merge UPCs with new heavy-ion machinery (Angantyr) recently introduced to PYTHIA 8
- Improve efficiency (currently optimized for ep)
- Hard diffraction for photoproduction (see I.H. on Friday)
 - Based on diffractive PDFs and dynamical rapidity gap survival from MPIs, extend to nuclear target
- Smooth merging of photoproduction and DIS

Backup slides

MPI and parton shower generation

Common evolution scale (p_T) for FSR, ISR and MPIs

- Probability for something to happen at given p_T

$$\frac{d\mathcal{P}}{dp_T} = \left(\frac{d\mathcal{P}_{\text{MPI}}}{dp_T} + \sum \frac{d\mathcal{P}_{\text{ISR}}}{dp_T} + \sum \frac{d\mathcal{P}_{\text{FSR}}}{dp_T} \right) \\ \times \exp \left[- \int_{p_T}^{p_T^{\text{max}}} dp'_T \left(\frac{d\mathcal{P}_{\text{MPI}}}{dp'_T} + \sum \frac{d\mathcal{P}_{\text{ISR}}}{dp'_T} + \sum \frac{d\mathcal{P}_{\text{FSR}}}{dp'_T} \right) \right]$$

where $\exp[...]$ is a Sudakov factor

(probability that nothing else has happened before p_T)

Simultaneous partonic evolution

1. Start the evolution from a scale related to the hard process
2. Sample p_T values for each \mathcal{P}_i , pick one with highest p_T
3. Continue from the sampled p_T until reach $p_{T\text{min}} \sim \Lambda_{\text{QCD}}$

Partonic evolution for resolved photons

DGLAP equations for photons

- Additional term due to $\gamma \rightarrow q\bar{q}$ splittings

$$\frac{\partial f_i^\gamma(x, Q^2)}{\partial \log(Q^2)} = \frac{\alpha_{\text{em}}}{2\pi} e_i^2 P_{i\gamma}(x) + \frac{\alpha_s(Q^2)}{2\pi} \sum_j \int_x^1 \frac{dz}{z} P_{ij}(z) f_j(x/z, Q^2)$$

where $P_{i\gamma}(x) = 3(x^2 + (1-x)^2)$ for quarks, 0 for gluons (LO)

- Solution has two components:



$$f_i^\gamma(x, Q^2) = f_i^{\gamma, \text{pl}}(x, Q^2) + f_i^{\gamma, \text{had}}(x, Q^2)$$

- Point-like part from perturbative QCD
- Non-perturbative input required for the hadron-like part

$$f_i^{\gamma, \text{had}}(x, Q_0^2) = N_i x^{a_i} (1-x)^{b_i}$$

Parameter fixed in a global analysis



Charged particle η dependence in ep collisions at HERA



[Eur.Phys.J. C10 (1999) 363-372]

Dijet in ep collisions at HERA

Pseudorapidity dependence of dijets [Eur.Phys.J. C23 (2002) 615-631]



- Good agreement with the data
- Some sensitivity to MPIs with $x_\gamma^{obs} < 0.75$